INFLAMMATORY BOWEL DISEASE SPECIALTY CARE PROGRAM



Phone: 617-254-5900

		PRESCRIBER INFORMATION:				
			ddress:			
	State: Zip:		ity:	State:	Zip:	
Phone: Alt. Phone:			Phone: Fax:		170	
Email:			 PI:			
	ender: O M O F Caregiver:		ax I.D.:			
Height: Weight: Allergies: 0						
3 STATEMENT	OF MEDICAL NECESSITY: (F	Please Attac	ch All Medical Documentation	n)		
Date of Diagnosis: Crohn's Disease Ulcerative Colitis Irritable Bowel Syndrome		Prior Indicate D Failed Treatments: and Length o		Orug Name of Treatment:		
			☐ Biologics			
Other: Serious or active infection present?						
Hep B ruled out or treatment started?			☐ Corticosteroids			
			☐ Immunosuppressants			
			☐ Methotrexate			
If Prior Authorization is Denied: ☐ Automatically Draft Appeal for Review ☐ Send Preferred Formulary Alternatives			□ Surgery _			
			Other _			
	N INFORMATION: (Please be	sure to cho		ntenance dose v		
Medication	Dosage & Strength	□ Industic	Direction	1 14 and 00	6	Refills 0
□ CIMZIA®	☐ Prefilled Syringe Starter Kit☐ 200mg/ml Prefilled Syringe☐ 200mg Lyophilized Powder		Induction Dose: Inject 400mg SC on day 1, 14 and 28 Maintenance: Inject 400mg SC every 4 weeks			0
□ HUMIRA®	☐ Crohn's Disease/ Ulcerative Colitis Starter Kit		Induction Dose: Inject 160mg SC on day 1, then 80mg SC on day 15, then switch to maintenance dose			0
	☐ 40mg/0.8ml Pen ☐ 40mg/0.8ml Prefilled Syringe					
			RA Complete form	0.400		
☐ SIMPONI®	□ 100mg/ml Smartject® Autoinjector	at week	□ Induction Dose: Inject 200mg SC at week 0, 100mg SC at week 2 and then switch to maintenance dose			0
	☐ 100mg/ml Prefilled Syringe		Maintenance: Inject 100mg SC every 4 weeks Induction Dose: Patient Weight <55kg: 260mg;			
	☐ 130mg/26ml Vial		n Dose: Patient Weight <55kg: 26 85kg: 390mg; >85 kg: 520mg adr			0
☐ STELARA®	□ 45mg/0.5ml Prefilled Syringe□ 90mg/ml Prefilled Syringe□ 45mg/0.5ml Vial	☐ Maintenance Dose: Inject 90mg SC 8 weeks after the initial intravenous dose, then every 8 weeks thereafter			1	
☐ UCERIS®	☐ 9mg Tablets	☐ Take one	tablet daily in the morning with or	without food	30	1
☐ XIFAXAN®	☐ 550mg Tablets	☐ Take one	tablet three times daily for 14 day	'S	42	
	•		,			
5 INJECTION T	RAINING: O Pharmacist to Prov	ide Training	O Patient Trained in MD Office	e O Manufactur	er Nurse	Support
		O Physician				
7 INSURANCE	INFORMATION: Please Include	Front and E	Back Copies of Pharmacy and	Medical Card		
8 PRESCRIBER	R SIGNATURE: I authorize pharmacy to act	as my designee for in	itiating and coordinating insurance prior authorization	ns, nursing services and patie	ent assistance p	orograms.
	Date: ubstitution Permitted		Signature:	Written	Date:	
First authorization approval and insurance be	nefits will be determined by the payor based upon the patient's eligibility, m	edical necessity, and the te	articlipation	in this program is not a guarantee of	prior authorization	or or payment.