

Call: 617-254-5900

## **INFLAMMATORY BOWEL DISEASE SPECIALTY CARE PROGRAM**

1 PATIENT INFORMATION:			2 PRESCRIBER INFORMATION:			
Name:			Name:			
Address:						
	State: Zip:					
		NPI: DEA:				
			Tax I.D.:			
Height: Wei	ght: Allergies:		Office Contact: Phone:			
3 STATEMENT	OF MEDICAL NECESSITY: (	(Please Atta	ach All Medical Documentation	า)		
Date of Diagnosis:			Prior Indicate Drug			
☐ Crohn's Disease ☐ Ulcerative Colitis ☐ Irritable Bowel Syndrome ICD-10:			Failed Treatments: and Length of Treatment:  □ 5-ASA			
Other:			☐ Biologics			
Serious or active infection present?			☐ Corticosteroids			
Hep B ruled out or treatment started? ☐ Yes ☐ No						
TB Test: ☐ Positive ☐ Negative Date:			☐ Immunosuppressants			
If Prior Authorization is Denied:  ☐ Automatically Draft Appeal for Review ☐ Send Preferred Formulary Alternatives			☐ Methotrexate			
			☐ Surgery			
			☐ Other			
4 PRESCRIPTIO	N INFORMATION: (Please be	e sure to ch	oose both induction and main	itenance dose w	here ap	plicable)
Medication	Dosage & Strength		Direction		QTY	Refills
□ CIMZIA®	☐ Prefilled Syringe Starter Kit	□ Inducti	on Dose: Inject 400mg SC on day 1	, 14 and 28	6	0
	☐ 200mg/ml Prefilled Syringe☐ 200mg Lyophilized Powder	_	☐ Maintenance: Inject 400mg SC every 4 weeks ☐			
☐ HUMIRA®	☐ Crohn's Disease/ Ulcerative Colitis Starter Kit		□ Induction Dose: Inject 160mg SC on day 1, then 80mg SC on day 15, then switch to maintenance dose			0
	☐ 40mg/0.8ml Pen ☐ 40mg/0.8ml Prefilled Syringe	☐ Maintenance: Inject 40mg SC every other week ☐			2	
	☐ Patient ha		<u> </u>			
☐ SIMPONI®	☐ 100mg/ml Smartject <sup>®</sup> Autoinjector	at weel	☐ Induction Dose: Inject 200mg SC at week 0, 100mg SC at week 2 and then switch to maintenance dose			0
	☐ 100mg/ml Prefilled Syringe		☐ Maintenance: Inject 100mg SC every 4 weeks			
☐ STELARA®	☐ 130mg/26ml Vial	☐ Induction Dose: Patient Weight <55kg: 260mg; >55kg to 85kg: 390mg; >85 kg: 520mg administered IV				0
	<ul><li>□ 45mg/0.5ml Prefilled Syringe</li><li>□ 90mg/ml Prefilled Syringe</li><li>□ 45mg/0.5ml Vial</li></ul>	☐ Maintenance Dose: Inject 90mg SC 8 weeks after the initial intravenous dose, then every 8 weeks thereafter			1	
☐ UCERIS®	☐ 9mg Tablets	☐ Take or	ne tablet daily in the morning with or	without food	30	1
☐ XIFAXAN®	☐ 550mg Tablets	☐ Take or	ne tablet three times daily for 14 days	s	42	
	<del>-</del>		<u>_</u>			
G INJECTION TO	RAINING: O Pharmacist to Pro	vide Training	Patient Trained in MD Office	O Manufacture	r Nurse	Support
	<b>ELIVERY:</b> O Patient's Home				1140100	опрроп
	NFORMATION: Please Include					
	SIGNATURE: I authorize pharmacy to ac				nt assistance	orograms.
Signature:			Signature:	D	ate:	
70000	nefits will be determined by the payor based upon the patient's eligibility,	medical necessity, and the			ior authorization	or of payment.