

Call: 617-254-5900

## **MULTIPLE SCLEROSIS SPECIALTY CARE PROGRAM**

1 PATIENT INFORMATION: Name:		2 PRESCRIBER INFORMATION:  Name:	
City:	State: Zip:	City: State: 2	Zip:
Phone:	Alt. Phone:	Phone: Fax:	
Email:		NPI: DEA:	
DOB: Gender: C	M OF Caregiver:	Tax I.D.:	
Height: Weight:	Allergies:	Office Contact: Phone:	
3 STATEMENT OF ME	DICAL NECESSITY: (Plea	se Attach All Medical Documentation)	
Date of Diagnosis:	ICD-10:	Relapse/Remitting	☐ Progressive
	t experienced a first clinical episode?	☐ Yes ☐ No ☐ Attach MRI Results Date:	
Does the patient have any contraince	lication(s) to therapy?   No Yes	If Yes:	
If Prior Authorization is Denied:	☐ Automatically Draft Appeal for Rev	iew Send Formulary Preferred Alternatives	
A PRESCRIPTION INFO	ORMATION: (Please be sur	e to choose both induction and maintenance dose wl	here applicable)
Medication	Dosage & Strength	Direction	QTY Refills
	☐ 30mca Prefilled Syringe	☐ Inject 30mcg IM once a week	
☐ AVONEX®	☐ 30mcg Single Dose Vial ☐ 30mcg Avonex Pen	☐ Titration: 7.5mcg weekly (over a 4 week period) until target dose is reached which is 30mcg	1 Kit
☐ BETASERON®	☐ 0.3mg Lyophilized Powder	☐ Inject 0.25mg (1ml) SC every other day	1 Kit
		☐ Titration: Weeks 1-2: Inject 0.0625mg/0.25ml SC every other day Weeks 3-4: Inject 0.125mg/0.50ml SC every other day Weeks 5-6: Inject 0.1875mg/0.75ml SC every other day Weeks 7 and onward: Inject 0.25mg/1ml SC every other day	
		······································	
☐ COPAXONE®	☐ 20mg Prefilled Syringe☐ 40mg Prefilled Syringe☐	☐ Inject 20mg SC daily ☐ Inject 40mg SC three times per week ☐ Other	1 Kit
☐ COPAXONE®	, ,	☐ Inject 20mg SC daily☐ Inject 40mg SC three times per week	1 Kit
☐ COPAXONE®	, ,	☐ Inject 20mg SC daily ☐ Inject 40mg SC three times per week ☐ Other	1 Kit
	□ 40mg Prefilled Syringe	□ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 5-6: 0.1875mg/0.75ml SC every other day	_
□ EXTAVIA®	□ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder	□ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 5-6: 0.1875mg/0.75ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day	_
□ EXTAVIA® □ GILENYA®	□ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule	□ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 5-6: 0.1875mg/0.75ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day □ Take one capsule by mouth once daily □ Other	1 Kit
□ EXTAVIA® □ GILENYA®	□ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule □ 20mg Prefilled Syringe	□ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 5-6: 0.1875mg/0.75ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day □ Take one capsule by mouth once daily □ Other □ Inject 20mg SC daily □ Titration Pack Rebidose (six 8.8mcg pre-filled autoinjectors and	1 Kit
☐ EXTAVIA® ☐ GILENYA® ☐ GLATOPA™	□ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule □ 20mg Prefilled Syringe □ Titration Pack □ 22mcg Prefilled Syringe □ 44mcg Prefilled Syringe □ 44mcg Prefilled Syringe □ Rebidose® 22mcg Autoinjector	□ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day □ Take one capsule by mouth once daily □ Other □ Inject 20mg SC daily □ Titration Pack Rebidose (six 8.8mcg pre-filled autoinjectors and six 22 mcgpre-filled autoinjectors) □ For 22mcg SC 3 times per week maintenance dose: • Weeks 1 & 2: Inject 4.4mcg 3 times per week • Weeks 3 & 4: Inject 11mcg 3 times per week • Weeks 5 and onward: Inject 22mcg 3 times per week □ For 44mcg SC 3 times per week maintenance dose: • Weeks 1 & 2: Inject 8.8mcg 3 times per week • Weeks 3 & 4: Inject 22mcg 3 times per week • Weeks 3 & 4: Inject 22mcg 3 times per week	1 Kit
□ EXTAVIA® □ GILENYA® □ GLATOPA™ □ REBIF®	□ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule □ 20mg Prefilled Syringe □ Titration Pack □ 22mcg Prefilled Syringe □ 44mcg Prefilled Syringe □ Rebidose® 22mcg Autoinjector □ Rebidose® 44mcg Autoinjector	□ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day □ Take one capsule by mouth once daily □ Other □ Inject 20mg SC daily □ Titration Pack Rebidose (six 8.8mcg pre-filled autoinjectors and six 22 mcgpre-filled autoinjectors) □ For 22mcg SC 3 times per week maintenance dose: • Weeks 1 & 2: Inject 4.4mcg 3 times per week • Weeks 3 & 4: Inject 11mcg 3 times per week • Weeks 5 and onward: Inject 22mcg 3 times per week □ For 44mcg SC 3 times per week maintenance dose: • Weeks 1 & 2: Inject 8.8mcg 3 times per week • Weeks 3 & 4: Inject 22mcg 3 times per week • Weeks 3 & 4: Inject 22mcg 3 times per week	1 Kit
□ EXTAVIA® □ GILENYA® □ GLATOPA™ □ REBIF® □ INJECTION TRAININ	□ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule □ 20mg Prefilled Syringe □ Titration Pack □ 22mcg Prefilled Syringe □ 44mcg Prefilled Syringe □ Rebidose® 22mcg Autoinjector □ Rebidose® 44mcg Autoinjector	□ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day □ Take one capsule by mouth once daily □ Other □ Inject 20mg SC daily □ Titration Pack Rebidose (six 8.8mcg pre-filled autoinjectors and six 22 mcgpre-filled autoinjectors) □ For 22mcg SC 3 times per week maintenance dose: • Weeks 1 & 2: Inject 4.4mcg 3 times per week • Weeks 3 & 4: Inject 11mcg 3 times per week • Weeks 5 and onward: Inject 22mcg 3 times per week • Weeks 1 & 2: Inject 8.8mcg 3 times per week • Weeks 3 & 4: Inject 22mcg 3 times per week • Weeks 3 & 4: Inject 22mcg 3 times per week • Weeks 5 and onward: Inject 44mcg 3 times per week • Weeks 5 and onward: Inject 44mcg 3 times per week	1 Kit
□ EXTAVIA® □ GILENYA® □ GLATOPA™ □ REBIF® □ INJECTION TRAINING PRODUCT DELIVER	□ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule □ 20mg Prefilled Syringe □ Titration Pack □ 22mcg Prefilled Syringe □ 44mcg Prefilled Syringe □ Rebidose® 22mcg Autoinjector □ Rebidose® 44mcg Autoinjector	□ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.50ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day □ Take one capsule by mouth once daily □ Other □ Inject 20mg SC daily □ Titration Pack Rebidose (six 8.8mcg pre-filled autoinjectors and six 22 mcgpre-filled autoinjectors) □ For 22mcg SC 3 times per week maintenance dose: • Weeks 1 & 2: Inject 4.4mcg 3 times per week • Weeks 3 & 4: Inject 11mcg 3 times per week • Weeks 5 and onward: Inject 22mcg 3 times per week • Weeks 1 & 2: Inject 8.8mcg 3 times per week • Weeks 3 & 4: Inject 22mcg 3 times per week • Weeks 3 & 4: Inject 22mcg 3 times per week • Weeks 5 and onward: Inject 44mcg 3 times per week • Weeks 5 and onward: Inject 44mcg 3 times per week	1 Kit
□ EXTAVIA® □ GILENYA® □ GLATOPA™ □ REBIF® □ MIJECTION TRAINING PRODUCT DELIVER 1 INSURANCE INFOR	□ 40mg Prefilled Syringe □ 0.3mg Lyophilized Powder □ 0.5mg Capsule □ 20mg Prefilled Syringe □ Titration Pack □ 22mcg Prefilled Syringe □ 44mcg Prefilled Syringe □ Rebidose® 22mcg Autoinjector □ Rebidose® 44mcg Autoinjector □ Rebidose® 44mcg Autoinjector □ Rebidose® Horizet Provide The Please Include From The Please	□ Inject 20mg SC daily □ Inject 40mg SC three times per week □ Other □ Inject 0.25mg (1ml) SC every other day □ Titration: Weeks 1-2: 0.0625mg/0.25ml SC every other day Weeks 3-4: 0.125mg/0.75ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day Weeks 7 and onward: 0.25mg/1ml SC every other day Underson other □ Inject 20mg SC daily □ Titration Pack Rebidose (six 8.8mcg pre-filled autoinjectors and six 22 mcgpre-filled autoinjectors) □ For 22mcg SC 3 times per week maintenance dose: ■ Weeks 1 & 2: Inject 4.4mcg 3 times per week ■ Weeks 3 & 4: Inject 11mcg 3 times per week ■ Weeks 5 and onward: Inject 22mcg 3 times per week ■ Weeks 1 & 2: Inject 8.8mcg 3 times per week ■ Weeks 3 & 4: Inject 22mcg 3 times per week ■ Weeks 3 & 4: Inject 22mcg 3 times per week ■ Weeks 3 & 4: Inject 22mcg 3 times per week ■ Weeks 5 and onward: Inject 44mcg 3 times per week ■ Weeks 5 and onward: Inject 44mcg 3 times per week ■ Weeks 5 and onward: Inject 44mcg 3 times per week	1 Kit 30 1 Kit